

## Most common tumor locations

Mariana A. Phillips, MD

Tumor	Most Common Location
Angiosarcoma	Head and neck
Blue nevus, cellular	Buttocks and sacrococcygeal region
Blue nevus, common	Near the dorsa of hands and feet
Clear cell acanthoma	Shin, calf, thigh
Clear cell hidradenoma	Face, eyelids
Chondroid syringoma	Neck and cheek
Dermatofibroma	Legs
Dermatofibrosarcoma protuberans	Trunk > proximal extremities
Hibernoma	Neck, axilla, shoulder
Juvenile xanthogranuloma (JXG)	Head and neck; iris
Leiomyosarcoma	Extensor surface of extremities
Liposarcoma	Upper thigh > buttocks
Lymphangioma circumscriptum	Abd, axilla, tongue
Merkel cell	Face
Mucinous eccrine carcinoma	Eyelid > face > axilla/trunk
Tufted hemangioma	Neck or trunk
Sebaceous carcinoma	Periocular (increased risk of mets)
Sinusoidal hemangioma	Female: Trunk and breasts
Smooth muscle hamartoma	Lumbosacral region
Spindle cell lipoma	Back (post trunk), neck, shoulder
Spitz nevus	Head, lower extremities

### References

Bolognia J, Jorizzo J, Rapini R, et al. *Dermatology*. Mosby Publishing; 2003.  
 Odum R, James W, Berger T. *Andrews' Disease of the Skin*. Ninth Edition. W.B. Saunders Company; 2000.



Mariana A. Phillips, M.D., is currently a Mohs fellow at the Virginia Commonwealth University in Richmond, Virginia.