
# DermWorld directions in residency

### boards fodder

#### JAK inhibitors in dermatology

By Jay Agarwal, DO



A Publication of the Am



Jay Agarwal, DO, is a PGY-4 resident at Beaumont-Trenton in Michigan.

Generic name	Brand name	Route	Primary MOA	FDA-approved indications	Side effects and adverse events	Other
Ruxolitinib	Opzelura	Topical	JAK1/JAK2 inhibitor	<u>Dermatologic:</u> ⇔Mild-to- moderate AD ⇔Non- segmental vitiligo	Common: URI, UTI, urticaria, diarrhea, follicu- litis, application site acne, applica- tion site pruritis, HA, pyrexia Less common: Anemia, neutrope- nia, thrombocyto- penia, lipid eleva- tions, NMSC	Black box warnings Increased risk of: 1. Serious infections (including TB and herpes zoster) 2. Mortality 3. Malignancies (including lymphoma) 4. MACE 5. Thrombosis First FDA-approved prescription treatment for vitiligo

## boards fodder

## JAK inhibitors in dermatology

By Jay Agarwal, DO

Generic name	Brand name	Route	Primary MOA	FDA-approved indications	Side effects and adverse events	Other
Baricitinib	Olumiant	Oral	JAK1/JAK2 inhibitor	Dermatologic: ⇒Severe AA <u>Other:</u> ⇒RA ⇒COVID-19	Common: URI, nausea, herpes simplex, herpes zoster, HA, acne, lipid elevations, CPK increase, UTI, liver enzyme eleva- tions, folliculitis, genital Candida infections, ane- mia, neutropenia, abdominal pain, weight increase Less common: Hypersensitivity, GI perforation, lymphopenia, NMSC	Black box warnings Increased risk of: 1. Serious infections (including TB) 2. Mortality 3. Malignancies (includ- ing lymphoma) 4. MACE 5. Thrombosis Avoid the use of live vaccines while on treat- ment
Ruxolitinib	Jakafi	Oral	JAK1/JAK2 inhibitor	<u>Dermatologic:</u> ⇔GVHD <u>Other:</u> ⇔Myelofibrosis ⇔PV	Common Anemia, neutro- penia, thrombocy- topenia, infection, edema Less common Serious infec- tion, NMSC, lipid elevations, MACE, thrombosis, sec- ondary malignan- cies	Avoid concomitant use with fluconazole doses greater than 200mg
Tofacitinib	Xeljanz	Oral	JAK1/3 inhibitor	Dermatologic: None <u>Other:</u> ⇒RA ⇒PsA ⇒UC ⇒AS ⇒JIA	Common: URI, diarrhea, HA, herpes zoster, lipid elevations, transa- minitis Less common: GI perforation, NMSC, pancyto- penia	Black box warnings Increased risk of: 1. Serious infections (including TB) 2. Mortality 3. Malignancies (includ- ing lymphoma) 4. MACE 5. Thrombosis Avoid the use of live vaccines while on treat- ment
Upadacitinib	Rinvoq	Oral	JAK1 inhibitor	Dermatologic: ⇒Moderate-to- severe AD <u>Other:</u> ⇒RA ⇒PsA ⇒UC ⇒AS	Common: URI, acne, herpes simplex, herpes zoster, influenza, HA, CPK increase, cough, folliculitis, abdominal pain, weight gain, neu- tropenia, myalgia, pyrexia Less common: Hypersensitivity, GI perforations, neutropenia, ane- mia, lymphopenia, increased LDL and HDL	Black box warnings   Increased risk of:   1. Serious infections   (including TB)   2. Mortality   3. Malignancies (including lymphoma)   4. MACE   5. Thrombosis   Avoid the use of live   vaccines while on treatment   Females of reproductive potential should be advised to use effective   contraception during   treatment and for 4   weeks following completion of therapy

## boards fodder

#### JAK inhibitors in dermatology

By Jay Agarwal, DO

Generic name	Brand name	Route	Primary MOA	FDA-approved indications	Side effects and adverse events	Other
Abrocitinib	Cibinqo	Oral	JAK1 inhibitor	<u>Dermatologic:</u> ⇔Moderate-to- severe AD -	Common: Nasopharyngitis, nausea, HA, CPK increase, UTI, acne, vomiting, oropharyngeal pain, influenza, gastroenteritis, impetigo, HTN, contact dermatitis, herpes simplex, herpes zoster, thrombocytopenia Less common: Lymphopenia, lipid elevations, retinal detachment	Black box warnings Increased risk of: 1. Serious infections (including TB) 2. Mortality 3. Malignancies (includ- ing lymphoma) 4. MACE 5. Thrombosis Avoid the use of live vaccines prior to, dur- ing, and immediately after treatment Contraindications: antiplatelet therapies except for low-dose aspirin (≤81 mg daily), during the first 3 months of treatment
Deucravacitinib	Sotyktu	Oral	TYK2 inhibitor	Dermatologic: ⇒Moderate-to- severe plaque psoriasis	Common: URI, CPK increase, herpes simplex, mouth ulcers, fol- liculitis, acne Less common: Hypersensitivity, herpes zoster, serious infections (including TB), malignancy (includ- ing lymphoma), rhabdomyolysis, hypertriglyceride- mia, transaminitis	No black box warning Avoid the use of live vaccines while on treat- ment

#### **Abbreviations**

AA = Alopecia areata

AD = Atopic dermatitis

AS = Ankylosing spondylitis

GVHD = Graft-versus-host disease

IBD = Inflammatory bowel disease

JIA = Juvenile idiopathic arthritis

MACE = Major adverse cardiac events (defined as cardiovascular death, myocardial infarction, and stroke)

PsA = Psoriatic arthritis PV = Polycythemia vera

RA = Rheumatoid arthritis

UC = Ulcerative colitis

#### UC = Ulcerative coll

#### Sources

www.opzelura.com/prescribing-information.pdf www.jakafi.com/pdf/prescribing-information.pdf https://pi.lilly.com/us/olumiant-uspi.pdf www.ema.europa.eu/en/documents/product-information/olumiant-epar-product-information\_en.pdf https://labeling.pfizer.com/ShowLabeling.aspx?id=959 www.rxabbvie.com/pdf/rinvoq\_pi.pdf www.accessdata.fda.gov/drugsatfda\_docs/label/2022/213871s000lbl.pdf https://packageinserts.bms.com/pi/pi\_sotyktu.pdf