DOCUMENTATION TIPS

MIPS #410: Psoriasis: **Clinical Response to Systemic Medications**

Electronic health records (EHRs) collect and organize notes, medication lists, and patient information using various formats. With providers also documenting this information in unique ways, this can potentially cause confusion and an increased timeline for measure mapping with DataDerm. This tip sheet can help you manage reporting requirements for performance measures and streamline standard documentation practices to allow seamless data pull into DataDerm.

The DataDerm team will work with you to connect DataDerm with your EHR to extract data. To make that as smooth as possible, it helps to document key elements of patient care. DataDerm cannot read scanned images of any kind, including scanned images for labs, letters to physicians, pathology reports, follow-up plans, and dates. If you have scanned images with information needed for your measures, please add a note in your chart with the date and required patient information for this data to be accurately collected.

This tip sheet can assist paper-based practices in standardizing documentation practices. Keeping notes in the patient's paper chart of all documentation requirements will assist you when reporting for this measure.

MIPS #410: Psoriasis: Clinical Response to Systemic Medications

For all patients, document the following in your notes:

- Current diagnosis of **psoriasis vulgaris**
 - Include the ICD-10 code L40.0 and
 - CPT or HCPCS codes for the patient encounter during measurement period (99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99324, 99325, 99326, 99327, 99328, 99334, 99335, 99336, 99337, 99341, 99342, 99343, 99344, 99345, 99347, 99348, 99349, 99350, G0438, G0439)
- Code G9764 Patient is treated with a systemic medication for psoriasis vulgaris
 - List systemic medication on medication list and/or identify systemics prescribed for treatment of psoriasis vulgaris (see medication list below)
 - Provider name that ordered the systemic medication in notes or identify systemics ordered by another provider
 - State date the systemic was prescribed
 - Document dates of any gap in medication use
 - Document if patient had a gap of 4 weeks or more during the course of treatment
- Assessment scale name, date provided, and results/score
 - Physician Global Assessment (PGA; 5-point or 6-point scale)
 - Body Surface Area (BSA)
 - Psoriasis Area and Severity Index (PASI)
 - Dermatology Life Quality Index (DLQI)
- Document if applicable:
 - Patient declined therapy change
 - Patient has documented contraindications
 - Patient has not been treated with a systemic for at least 6 consecutive months

Page 1 of 3

For more information, contact the American Academy of Dermatology:

WEBSITE: aad.org



 $^{\odot}$ by the American Academy of Dermatology and the American Academy of Dermatology Association. "All Right's Reserved." Reproduction or republication is strictly prohibited without prior written permission.

DOCUMENTATION TIPS

MIPS #410: Psoriasis: Clinical Response to Systemic Medications

MIPS #410: Psoriasis: Clinical Response to Systemic Medications

Additional Tips:

- Collect **once per performance period** for patients seen during the performance period.
- This measure **ONLY** applies to patients with a diagnosis of **psoriasis vulgaris (L40.0)**.
- See measure specifications for additional codes that can assist in seamless measure mapping from your EHR to DataDerm, if applicable (e.g. G9649).
- GQ, GT, 95, and POS 02 telehealth modifiers make cases ineligible.
- Go to the DataDerm Dashboard Resources" tab to get the assessment tools for this measure. Click "Tools" for resources on picking the best assessment tool for your workflow.

Examples of Applicable Medications for MIPS 410

| Brand name | Chemical name | Mechanism of action/ Type of biologic | Indications |
|------------|-----------------|--|---|
| Stelara | Ustekinumab | Interleukin (IL) 12 and 23 inhibitor | Psoriasis and/or psoriatic arthritis |
| Cosentyx | Secukinumab | IL-17 inhibitor | Psoriasis and/or psoriatic arthritis |
| Siliq | Brodalumab | IL-17 inhibitor | Psoriasis |
| Taltz | lxekizumab | IL-17 inhibitor | Psoriasis and/or psoriatic arthritis |
| Tremfya | Guselkumab | IL-23 inhibitor | Psoriasis |
| Amjevita | Adalimumab-atto | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Cimzia | Certolizumab | TNF-alpha inhibitor | Psoriatic arthritis |
| Cyltezo | Adalimumab-adbm | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Enbrel | Etanercept | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Erelzi | Etanercept-szzs | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |

Page 2 of 3

For more information, contact the American Academy of Dermatology: WEBSITE: aad.org

© by the American Academy of Dermatology and the American Academy of Dermatology Association. "All Rights Reserved." Reproduction or republication is strictly prohibited without prior written permission.

The Measure is copyrighted but can be reproduced and distributed, without modification, for noncommercial purposes (eg, use by healthcare providers in connection with their practices) without obtaining approval from NCQA.



DOCUMENTATION TIPS

MIPS #410: Psoriasis: Clinical Response to Systemic Medications

| Brand name | Chemical name | Mechanism of action/ Type of biologic | Indications |
|-------------------------|-----------------------|---|---|
| Humira | Adalimumab | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Inflectra | Infliximab-dyyb | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Remicade | Infliximab | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Renflexis | Infliximab-abda | TNF-alpha inhibitor | Psoriasis and/or psoriatic arthritis |
| Simponi & Simponia Aria | Golimumab | TNF-alpha inhibitor | Psoriatic arthritis |
| Rheumatrex | Methotrexate | Blocks dihydrofolate reductase | Psoriasis |
| Trexall | Methotrexate | Blocks dihydrofolate reductase | Psoriasis |
| Xatmep | Methotrexate | Blocks dihydrofolate reductase | Psoriasis |
| Otrexup | Methotrexate | Inhibits dihydrofolic acid reductase | Psoriasis |
| Rasuvo | Methotrexate | Inhibits dihydrofolic acid reductase | Psoriasis |
| Gengraf | Cycloporine, modified | Blocks lymphocytes (T cells) and lymphokines | Psoriasis |
| Neoral | Cycloporine, modified | Blocks lymphocytes (T cells) and lymphokines | Psoriasis |
| Sandimmune | Cyclosporine | Blocks lymphocytes (T cells) and lymphokines | Psoriasis |
| Otezla | Apremilast | Blocks phosphodiesterase-4 (PDE4) | Psoriasis and/or psoriatic arthritis |
| Soriatane | Acitretin | Unknown; activates retinoid receptors | Psoriasis |
| llumya | Tildrakizumab | Selective (IL)-23p19 inhibitor | Psoriasis |

Page 3 of 3

For more information, contact the American Academy of Dermatology: WEBSITE: aad.org

© by the American Academy of Dermatology and the American Academy of Dermatology Association. "All Rights Reserved." Reproduction or republication is strictly prohibited without prior written permission.

The Measure is copyrighted but can be reproduced and distributed, without modification, for noncommercial purposes (eg, use by healthcare providers in connection with their practices) without obtaining approval from NCQA.

